
```

<?xml version="1.0" encoding="UTF-8"?>
<!DOCTYPE simulation [
<!ENTITY Npts      "64">
<!ENTITY Nsamples  "64">
<!ENTITY L         "3e-5">
]>
<simulation xmzs-version="2">
  <name>GPE_1D_groundstate_course</name>

  <author> Sebastian Wuester </author>
  <description>
    Finds ground state of Gross-Pitaevskii-equation in 1D, SI units
  </description>

  <geometry>
    <propagation_dimension> t </propagation_dimension>
    <transverse_dimensions>
      <dimension name="x" lattice="&Npts;" domain="(-&L;, &L;)" />
    </transverse_dimensions>
  </geometry>

  <features>
    <benchmark />
    <auto_vectorise />
    <fftw />
    <globals>
      <![CDATA[
const double hbar = 1.05457266e-34;
const double omega = 10.0*(2.0*M_PI);
const double omega_perp = 200.0*(2.0*M_PI);
const double x0 = 0.0;

// Rb 87
const double mass = 1.4432e-25;
const double as = 5.5e-9;

//dervied quantities
const double Natoms = 100.0;

const double U = 4.0*M_PI*hbar*hbar*as/mass;
const double sigma = sqrt(hbar/mass/omega);
const double sigma_perp = sqrt(hbar/mass/omega_perp);
const double U1d = U/(2.0*M_PI*sigma_perp*sigma_perp);
const double normfact = pow(M_PI*sigma*sigma,-0.25);

]]>
    </globals>
  </features>

  <vector name="wavefunction" initial_space="x" type="complex">
    <components>psi</components>
    <initialisation>
      <![CDATA[
const double delx = x - x0;
psi = normfact*sqrt(Natoms)*exp(-0.5*delx*delx/sigma/sigma);
]]>
    </initialisation>
  </vector>

  <vector name="potentials" initial_space="x" type="real">
    <components>trap</components>

```

```
<initialisation>
  <![CDATA[
    trap=0.5*mass*omega*omega*x*x;
  ]]>
</initialisation>
</vector>

<computed_vector name="moments" dimensions="" type="real">
  <components> norm expecxx expecx </components>
  <evaluation>
    <dependencies basis="x"> wavefunction </dependencies>
    <![CDATA[
      norm = mod2(psi);
      expecxx = x*x*mod2(psi);
      expecx = x*mod2(psi);
    ]]>
  </evaluation>
</computed_vector>

<sequence>
  <integrate algorithm="RK4" interval="5.0" steps="1000000">
    <samples>200 200 200</samples>
    <!-- -->
    <filters where="step end">
      <filter>
        <dependencies>wavefunction moments</dependencies>
        <![CDATA[
          // Correct normalisation of the wavefunction
          psi *= sqrt(Natoms/norm);
        ]]>
      </filter>
    </filters>
    <!-- -->
    <operators>
      <operator constant="yes" kind="ip">
        <operator_names>L</operator_names>
        <![CDATA[
          L = -0.5*hbar*kx*kx/mass;
        ]]>
      </operator>
      <integration_vectors>wavefunction</integration_vectors>
      <dependencies>potentials</dependencies>
      <![CDATA[
        double dens=psi.Re()*psi.Re() + psi.Im()*psi.Im();

        dpsi_dt = L[psi] - (Uld*dens + trap)*psi/hbar;
      ]]>
    </operators>
  </integrate>
  <!-- -->
  <breakpoint filename="groundstate_break" format="hdf5">
    <dependencies basis="x"> wavefunction </dependencies>
  </breakpoint>
  <!-- -->
</sequence>

<output format="hdf5">
  <group>
    <sampling basis="x(&Nsamples;)" initial_sample="yes">
      <moments>density psire psiiim trappotential interaction_term</moments>
      <dependencies>wavefunction potentials</dependencies>
```

```
<![CDATA[
    density = mod2(psi);
    psire = psi.Re();
    psiim = psi.Im();
    trappotential = trap;
    interaction_term = Uld*mod2(psi);
]]>
</sampling>
</group>
<group>
    <sampling basis="kx(&Npts;)" initial_sample="yes">
        <moments>fspec</moments>
        <dependencies>wavefunction</dependencies>
        <![CDATA[
            fspec = mod2(psi);
        ]]>
    </sampling>
</group>
<group>
    <sampling basis="" initial_sample="yes">
        <moments> atomnumber meanpos deltapos </moments>
        <dependencies> moments </dependencies>
        <![CDATA[
            atomnumber = norm;
            meanpos = expecx/norm;
            deltapos = sqrt(expecxx/norm - (expecx/norm)*(expecx/norm));
        ]]>
    </sampling>
</group>
</output>
</simulation>
```