

```
<?xml version="1.0" encoding="UTF-8"?>
<!DOCTYPE simulation [
<!ENTITY Npts      "64">
<!ENTITY Nsamples   "64">
<!ENTITY L         "3e-5">
]>
<simulation xmds-version="2">
  <name>GPE_1D_noisy_course_MT</name>

  <author> Sebastian Wuester </author>
  <description>
    Gross-Pitaevskii-equation in 1D, SI units
  </description>

  <geometry>
    <propagation_dimension> t </propagation_dimension>
    <transverse_dimensions>
      <dimension name="x" lattice="&Npts;" domain="(-&L;, &L;)" />
    </transverse_dimensions>
  </geometry>

  <features>
    <benchmark />
    <auto_vectorise />
    <fftw />
    <globals>
      <![CDATA[
        const double hbar = 1.05457266e-34;
        const double omega = 10.0*(2.0*M_PI);
        const double omega_perp = 200.0*(2.0*M_PI);

        // Rb 87
        const double mass = 1.4432e-25;
        const double as = 5.5e-9;

        const double noiseamp = 0.2;

        //dervied quantities
        const double Natoms = 100.0;

        const double U = 4.0*M_PI*hbar*hbar*as/mass;
        const double sigma = sqrt(hbar/mass/omega);
        const double sigma_perp = sqrt(hbar/mass/omega_perp);
        const double U1d = U/(2.0*M_PI*sigma_perp*sigma_perp);
        const double normfact = pow(M_PI*sigma*sigma,-0.25);

      ]]>
    </globals>
  </features>

  <driver name="multi-path" paths="100" />

  <vector name="wavefunction" initial_space="x" type="complex">
    <components>psi</components>
    <initialisation kind="hdf5">
      <filename> groundstate_break.h5</filename>
    </initialisation>
  </vector>

  <vector name="potentials" initial_space="x" type="real">
    <components>trap</components>
```

```
<initialisation>
  <![CDATA[
    trap=0.5*mass*omega*omega*x*x;
  ]]>
</initialisation>
</vector>

<computed_vector name="moments" dimensions="" type="real">
  <components> norm expecxx expecx </components>
  <evaluation>
    <dependencies basis="x"> wavefunction </dependencies>
    <![CDATA[
      norm = mod2(psi);
      expecxx = x*x*mod2(psi);
      expecx = x*mod2(psi);
    ]]>
  </evaluation>
</computed_vector>

<noise_vector name="thermalNoise" dimensions="x" kind="Gaussian"
               type="complex" method="dsfmt" seed="314 159 276">
  <components>Eta</components>
</noise_vector>

<sequence>
  <!-- -->
  <filter>
    <dependencies>wavefunction thermalNoise</dependencies>
    <![CDATA[
      psi += noiseamp*Eta;
    ]]>
  </filter>
  <!-- -->
  <integrate algorithm="ARK89" interval="0.4" tolerance="1e-8">
    <samples>200 200 200</samples>
    <operators>
      <operator kind="ip">
        <operator_names>L</operator_names>
        <![CDATA[
          L = -i*0.5*hbar*kx*kx/mass;
        ]]>
      </operator>
      <integration_vectors>wavefunction</integration_vectors>
      <dependencies>potentials</dependencies>
      <![CDATA[
        double dens=psi.Re()*psi.Re() + psi.Im()*psi.Im();

        dpsi_dt = L[psi] - i*(U1d*dens + trap )*psi/hbar;
      ]]>
    </operators>
  </integrate>
</sequence>

<output format="hdf5">
  <group>
    <sampling basis="x(&Nsamples;)" initial_sample="yes">
      <moments>density psire psiim trappotential interaction_term</moments>
      <dependencies>wavefunction potentials</dependencies>
      <![CDATA[
        density = mod2(psi);
        psire = psi.Re();
      ]]>
    </sampling>
  </group>
</output>
```

```
psiim = psi.Im();
trappotential = trap;
interaction_term = U1d*mod2(psi);
]]>
</sampling>
</group>
<group>
<sampling basis="kx(&Npts;)" initial_sample="yes">
<moments>fspec</moments>
<dependencies>wavefunction</dependencies>
<![CDATA[
    fspec = mod2(psi);
]]>
</sampling>
</group>
<group>
<sampling basis="" initial_sample="yes">
<moments> atomnumber meanpos deltapos </moments>
<dependencies> moments </dependencies>
<![CDATA[
    atomnumber = norm;
    meanpos = expecx/norm;
    deltapos = sqrt(expecxx/norm - (expecx/norm)*(expecx/norm));
]]>
</sampling>
</group>
</output>
</simulation>
```