

Delocalization and spreading in a nonlinear Stark ladder

Dmitry O. Krimer,¹ Ramaz Khomeriki,^{1,2} and Sergej Flach¹

¹Max-Planck Institute for the Physics of Complex Systems, Nöthnitzer Str. 38, 01187 Dresden, Germany

²Physics Department, Tbilisi State University, Chavchavadze 3, 0128 Tbilisi, Georgia

(Received 18 April 2009; revised manuscript received 2 July 2009; published 4 September 2009)

We study the evolution of a wave packet in a nonlinear Stark ladder. In the absence of nonlinearity all normal modes are spatially localized giving rise to an equidistant eigenvalue spectrum and Bloch oscillations. Nonlinearity induces frequency shifts and mode-mode interactions and destroys localization. For large strength of nonlinearity we observe single-site trapping as a transient, with subsequent explosive spreading, followed by subdiffusion. For moderate nonlinearities an immediate subdiffusion takes place. Finally, for small nonlinearities we find linear Stark localization as a transient, with subsequent subdiffusion. For single-mode excitations and weak nonlinearities, stability intervals are predicted and observed upon variation in the dc bias strength, which affects the short- and the long-time dynamics.

DOI: [10.1103/PhysRevE.80.036201](https://doi.org/10.1103/PhysRevE.80.036201)

PACS number(s): 05.45.-a, 05.60.Cd, 63.20.Pw

The textbook solid-state problem of an electron in a periodic potential with an additional electric field (see, e.g., [1]) leads to investigations of Bloch oscillations [2] and Landau-Zener tunneling [3] in different physical systems. Optical waves in photonic lattices [4,5] and ultracold atoms in optical lattices [6–8] are recent examples. If the well depth of the periodic potential is large enough and the Landau-Zener tunneling is suppressed, the problem is discretized (Stark ladder) and the resulting eigenvalue problem is explicitly solved in terms of the localized eigenmodes [9] of the system. Additional nonlinear terms in the wave equations are generated, e.g., by a nonlinear response of the medium (light) or by a mean-field treatment of quantum many-body interactions (Bose-Einstein condensates). These terms destroy the integrability of the wave equations leading to decoherence and deterioration of Bloch oscillations, which was observed on time scales of a few oscillation periods [5,7]. The study of hybrid discrete solitons in [10] showed that nonlinear Bloch oscillations may destroy solitary excitations.

Single-site excitations were studied in [11] on rather short time scales. In particular, three different ranges of nonlinearity with different dynamics for one value of the dc bias have been identified. Here, we present a detailed study and explain the delocalization process induced by nonlinear terms when starting from both single-site and single-mode excitations. We obtain the complete map of dynamical regimes in the parameter space of nonlinearity and the strength of the dc bias. We find a surprisingly abrupt explosive dynamics of the initially trapped excitations for strong nonlinearity. It can be experimentally verified in systems dealing, e.g., with ultracold atoms in optical lattices [8] and light propagation in nonlinear waveguide networks [5]. We also demonstrate a very peculiar dynamics for single-mode excitations and small nonlinearities. Depending on the strength of the dc bias, an initially excited mode might pump one of the neighboring modes or keep its amplitude for rather long times. These results can be observed experimentally for light propagating in waveguide networks where initial states can be prepared with well-defined phases [12].

We consider a discrete nonlinear Schrödinger equation with a dc bias E ,

$$i\dot{\Psi}_n = -(\Psi_{n+1} + \Psi_{n-1}) + nE\Psi_n + \beta|\Psi_n|^2\Psi_n, \quad (1)$$

where the overdot denotes the time derivative, Ψ_n is a complex amplitude of a wave at lattice site n , and β is the nonlinearity strength. Varying β is strictly equivalent to varying the norm $\mathcal{N} = \sum_n |\Psi_n|^2$. We choose $\mathcal{N} = 1$.

For $\beta = 0$, Eq. (1) is reduced to an eigenvalue problem using the ansatz $\Psi_n(t) = \exp(-i\lambda t)A_n$. In the case of an infinite lattice the eigenvalues $\lambda_\nu = E\nu$ (with ν being an integer) form an equidistant spectrum that extends over the whole real axis—the Stark ladder. The corresponding eigenfunctions or normal modes (NMs) obey the generalized translational invariance $A_{n+\mu}^{(\nu+\mu)} = A_n^{(\nu)}$ [2]. With $J_{n+1}(x) + J_{n-1}(x) = (2n/x)J_n(x)$ for the Bessel function $J_n(x)$ of the first kind [13], it follows that [9]

$$A_n^{(0)} = J_n(2/E). \quad (2)$$

All NMs are spatially localized with an asymptotic decay $|A_{n \rightarrow \infty}^{(0)}| \rightarrow (1/E)^n/n!$, giving rise to the well-known Bloch os-




FIG. 1. (Color online) (a) Black solid line: localization volume \mathcal{L} of the eigenfunction $A_n^{(0)}$ versus E . Insets: explicit form of the eigenfunction across the chain for two values $E=2$ and $E=0.2$ (b, blue; r, red); (b) The diagram of the three regimes of spreading in the parameter space (β, E) . Empty and filled circles: numerically obtained thresholds which separate the three different regimes I–III. Lines connecting symbols guide the eye. Black dashed line: threshold between II and III obtained from the dimer model (see text). In the limit of $E \rightarrow \infty$ all lines merge to the asymptotic limit $\beta \propto E$.

cillations with period $T_B=2\pi/E$. In Fig. 1(a) we plot the localization volume $\mathcal{L}=1/\sum_n |A_n^{(0)}|^4$ of an eigenstate, which characterizes the spatial extent of the NM, as a function of E . We find the asymptotic behavior of $\mathcal{L} \propto -[E \ln E]^{-1}$ for $E \rightarrow 0$. Note that, for $E=10$, \mathcal{L} is already close to its asymptotic value $\mathcal{L}=1$ for $E \rightarrow \infty$. The eigenvalue spacing E and the eigenvalue variation over a localization volume $\Delta \equiv E\mathcal{L}$ determine the packet evolution in the presence of nonlinearity.

With

$$\Psi_n(t) = \sum_{\nu} \phi_{\nu}(t) A_n^{(\nu)} e^{-i\nu Et}, \quad (3)$$

Eq. (1) transforms into NM space

$$i\dot{\phi}_{\nu} = \beta \sum_{\nu_1, \nu_2, \nu_3} I_{\nu, \nu_1, \nu_2, \nu_3} \phi_{\nu_1}^* \phi_{\nu_2} \phi_{\nu_3} e^{i(\nu + \nu_1 - \nu_2 - \nu_3)Et}, \quad (4)$$

where

$$I_{\nu, \nu_1, \nu_2, \nu_3} \equiv \sum_n A_{n-\nu}^{(0)} A_{n-\nu_1}^{(0)} A_{n-\nu_2}^{(0)} A_{n-\nu_3}^{(0)} \quad (5)$$

are the overlap integrals between eigenfunctions (2). To characterize the wave-packet dynamics we compute (i) the participation number $P=1/\sum_{\nu} |\phi_{\nu}|^4$ which measures the number of strongest excited NMs, (ii) the second moment $m_2 = \sum_{\nu} (\nu - \bar{\nu})^2 |\phi_{\nu}|^2$ ($\bar{\nu} = \sum_{\nu} \nu |\phi_{\nu}|^2$) which quantifies the degree of spreading of the wave packet, and (iii) the compactness index $\zeta = P^2/m_2$ which measures the degree of sparsity of the wave packet.

First, we study a single-site initial excitation $\Psi_n(0) = \delta_{n0}$. In that case the amplitudes in NM space are $\phi_{\nu}(0) = J_{\nu}(2/E)$. The nonlinear frequency shift at site $n=0$ is $\delta \equiv \beta$. We expect three qualitatively different regimes of spreading: (I) $\delta < E$, (II) $E < \delta < \Delta$, and (III) $\Delta < \delta$. In case (I) the nonlinear frequency shift is less than the spacing between excited modes. Therefore, no initial resonance overlap is expected, and the dynamics may evolve as the one for $\beta=0$ at least for long times. In case (II) resonance overlap happens, and the packet should evolve differently. For case (III) δ tunes the excited site out of resonance with the neighboring NMs. Resonances with more distant NMs are possible, but the overlap with these NMs is the weaker the further away they are, see Eq. (2). Therefore, for long times the excited site may evolve as an independent oscillator (trapping). The diagram of the three regimes of spreading in the parameter space (β, E) is shown in Fig. 1(b).

Let us start with case (III). For $E=2$ and $\beta > 8.9$, the single-site excitation stays trapped up to times $t=10^8$ without significantly spreading into any other site of the lattice (violet curve in Fig. 2). Slightly lowering β we observe that the excitation is trapped up to a some time T_E which sensitively depends on β and changes by many orders of magnitude, e.g., between 10^2 and 10^7 in the narrow interval $\beta \in (8.05, 8.9)$ for $E=2$ (Fig. 2). For times $t > T_E$ an explosive and spatially asymmetric spreading is observed on a time scale of one Bloch period T_B . The packet spreads in the direction of NMs with larger eigenvalues, which provide the possibility of resonant energy transfer from the single-site excitation due to its positive nonlinear frequency shift δ . For




FIG. 2. (Color online) Single-site excitation for $E=2$. Second moment m_2 and participation number P versus time in log-log plots for different values of β inside the interval where an explosive delocalization of the trapped regime occurs: $\beta=8.15, 8.25, 8.5$ (o, orange; g, green; r, red). $\beta=8$ (b, blue): intermediate regime. $\beta=8.9$ (v, violet): trapped regime.

about ten Bloch periods T_B the packet shows Bloch oscillations, which then quickly decohere. Finally, the packet spreads incoherently and subdiffusively (Figs. 2 and 3). The explosion time T_E is not monotonously changing with β (cf. Fig. 2), which indicates intermittency, i.e., the single-site excitation can be closer or further away from some regular structures in phase space. That distance may in turn control the value of T_E . For $E=2$ and $\beta=8$ the packet spreads from scratch (blue curve in Fig. 2). Such dynamics can be experimentally verified in systems dealing, e.g., with ultracold atoms in optical lattices [8] and light propagation in nonlinear waveguide networks [5].

In regime II the nonlinearity is not strong enough to trap a single-site excitation, but the excited NMs are resonantly interacting. Therefore, the packet expands subdiffusively from the very beginning (green curve in Fig. 4). Since the eigenmodes of the linear system are spatially localized, the incoherent way of subdiffusive spreading resembles the one ob-




FIG. 3. (Color online) $|\Psi_n(t)|^2$ as a function of time (in logarithmic scale) and lattice site for $E=2, \beta=8.15$ (upper graph) and $E=0.2, \beta=5.2$ (lower graph). The zooms into the explosion region are plotted on a linear time scale. Bloch oscillations are observed right after the explosion time T_E .




FIG. 4. (Color online) Single-site excitation for $E=2$. m_2 and P versus time (log-log plots) for (i) $\beta=2.1$ (b, blue): weak nonlinear regime; (ii) $\beta=4$ (g, green): intermediate regime; and (iii) $\beta=9$ (r, red): trapped regime. Orange dashed line: $m_2 \sim t^{0.38}$. Inset: ζ for $\beta=4$.

served in disordered nonlinear systems without [14–16] and with [17] a dc bias. The second moment m_2 appears to grow without bounds according to $m_2 \sim t^\alpha$. We fitted the exponent α for 20 different runs for $E=2$ and $\beta=4$. Each run was obtained as follows. We integrated the equations up to $t=10^2$. The obtained distribution was transformed to NM space. The phases of all NMs were randomized, but the amplitudes were kept constant. Finally, we performed an inverse transformation to real space and continued the integration up to 10^8 . As a result we obtained the average exponent $\alpha=0.38 \pm 0.015$. Let us reconsider Eq. (4) and keep only resonant terms $\nu + \nu_1 - \nu_2 - \nu_3 = 0$. The resulting resonant normal form equations read

$$i\dot{\phi}_\nu = \beta \sum_{\nu_1, \nu_2} I_{\nu, \nu_1, \nu_2, (\nu + \nu_1 - \nu_2)} \phi_{\nu_1}^* \phi_{\nu_2} \phi_{\nu + \nu_1 - \nu_2}. \quad (6)$$

If at least two neighboring NMs are excited, the resonant normal form (6) will have a connectivity similar to the original lattice equations, which allows spreading over the whole lattice. Excluding the possibility of integrability of Eq. (6), we expect this spreading to happen indeed, as observed in the numerical runs. Following [14,15], we conclude that all NMs are in resonance with their neighborhood. If the wave packet is in a state of equipartition at any time, it follows that $\alpha=0.5$ [14,15]. In such an assumed equipartitioned state the compactness index $\zeta \approx 3$ [15]. However, the compactness index ζ decreases in time way beyond the equipartition threshold (inset in Fig. 4) and therefore the assumption of equipartitioned wave packets is incorrect. This is the reason for the discrepancy between the expected (0.5) and the observed (0.38) values of the exponent α . Note that the onset of subdiffusive spreading was also observed in Refs. [11,18] for two runs on rather short scales up to $t=10^5$, which makes it impossible to estimate the exponent.

In regime I the nonlinear frequency shift $\delta < E$. No strong initial resonance overlap between NMs is expected. In that case the dynamics may (at least for long times) evolve as in the linear case. Eventually, the subdiffusive spreading takes place after some waiting time τ (Fig. 4). Note that the cross-over from localization to spreading around time τ is not accompanied by a rapid redistribution of the energy between many neighboring sites as in the case of trapping with large nonlinearities. Therefore, the onset of subdiffusion is simply

related to the fact that weakly chaotic dynamics is characterized by a small momentary diffusion rate, and the corresponding diffusion process is observed only after times larger than τ .

In Fig. 1(b) we plot the boundaries between the three regimes in the parameter space (β, E) . These boundaries are determined as follows. Keeping the value of E fixed and varying β , we integrate the equations up to $t=10^5$ starting from the single-site excitation. Then we calculate m_2 and P of the final wave packet. Being in regime I, these values are quite close to those for the linear case. For a critical value of β they start to grow indicating that we enter regime II. Finally, at a second and larger value of β the values of m_2 and P drop down again, signaling the entering of regime III.

The border between regimes II and III can be accounted for by using a dimer model. Since the trapped state corresponds to only a single site being excited, we consider a dimer model where only one further lattice site (to the right) is added, in order to model the asymmetric energy transfer during the explosion. The resulting equations read

$$i\dot{\Psi}_0 = -\Psi_1 + \beta|\Psi_0|^2\Psi_0, \quad i\dot{\Psi}_1 = -\Psi_0 + E\Psi_1 + \beta|\Psi_1|^2\Psi_1,$$

which are solved using elliptic functions [19]. The dimer model is integrable, and the analytical computation of the initial condition $|\Psi_n(0)|^2 = \delta_{n,0}$ crossing a separatrix in the phase space of the dimer yields a relation between β and E . The transition to the trapped state at some critical nonlinearity $\beta(E)$ is accompanied by the divergence of the period of oscillations, and above the threshold the energy is mostly concentrated on one site. The final equation to be solved numerically reads

$$-\beta^3 E + \beta^2(3E^2 - 1) + \beta E(20 - 3E^2) + (E^2 + 4)^2 = 0. \quad (7)$$

As seen from Fig. 1(b) it fits very well with the boundary between regimes II and III obtained from the evaluation of the full set of equations. In particular, Eq. (7) correctly predicts that $\beta(E \rightarrow 0) \rightarrow 4$ and $\beta(E \rightarrow \infty) \rightarrow E$. Note that two critical nonlinearity values for self-trapping at $E=0.5, 5$ were found in [11], which agree reasonably with our data in Fig. 1(b).

We use the SBAB₂ symplectic integrator [20] to integrate Eq. (1) numerically. In order to ensure that our results are not generated by inaccurate computations, we performed different tests. In particular, we varied the time steps of our scheme by orders of magnitude. We also varied the size of the system to exclude finite-size effects in the evolution of the wave packets (finally, N was chosen to be from 500 to 1000). We found no visible change in the spreading characteristics. The narrow interval of nonlinearity with an explosive behavior remains also unchanged [21]. It is worth noting that the issue about numerical accuracy is discussed in details in the context of nonlinear disordered system in [15]. There we gave even more evidence that the round-off errors with double (or even single) precision are not affecting the results.

A single-mode excitation $\phi_\nu(t=0) = \delta_{\nu,0}$ also exhibits the three different regimes of spreading. However, for small values of nonlinearity β an intriguing feature of the short-time

dynamics follows. Indeed, consider the resonant normal form (6). For a single-mode excitation, it follows strictly that no other NM is going to be excited. Therefore, it is an exact solution of Eq. (6),

$$\phi_0(t) = \exp(-i\beta I_{0,0,0}t). \quad (8)$$

However, the full set of equations (4) will excite other NMs as well. These small perturbations may stay small or start to grow, depending on the stability of solution (8) of Eqs. (6). The linear equations for small perturbations with $\nu \neq 0$ read

$$i\dot{\phi}_\nu = \beta[I_{-\nu,\nu,0}\phi_0^2\phi_{-\nu}^* + 2I_{\nu,\nu,0}\phi_\nu], \quad (9)$$

$$i\dot{\phi}_{-\nu}^* = -\beta[I_{-\nu,\nu,0}\phi_0^2\phi_\nu + 2I_{\nu,\nu,0}\phi_{-\nu}^*].$$

Note that only pairs of NMs with $\pm\nu$ are coupled. With $\phi_{\pm\nu} \sim \exp(-i\beta I_{0,0,0}t + \gamma_\nu t)$, one gets

$$\gamma_\nu = |\beta| \sqrt{I_{-\nu,\nu,0}^2 - (2I_{\nu,\nu,0} - I_{0,0,0})^2}. \quad (10)$$

Thus, if $I_{-\nu,\nu,0}^2 > (2I_{\nu,\nu,0} - I_{0,0,0})^2$, it follows that γ_ν is real, the corresponding perturbations will grow exponentially fast in time, and the original single-mode excitation is unstable. Since the overlap integrals depend on the dc bias E , we expect transitions from stability to instability upon varying E . Indeed, a numerical evaluation of that condition yields that the single-mode excitation is stable in the limits $1.134 < E < 1.23$, $1.314 < E < 1.917$, and $E > 2.314$; otherwise, it is unstable. In the instability window $1.917 < E < 2.314$ ($1.23 < E < 1.314$) the modes $\nu = \pm 1$ ($\nu = \pm 2$) become unstable whereas the other modes remain stable. This behavior is confirmed by numerical simulations presented in Fig. 5 on short time scales. After sufficient exponential growth, the amplitude of the unstable mode saturates due to nonlinear corrections. Due to recurrences the energy eventually comes back to the initially excited mode after possible long times. We also find a correlation between the short-time stability properties of a single-mode excitation and the long-time evolution of the wave packet. If the single-mode excitation is launched in a stability window, the wave packet is practically not spreading up to $t = 10^8$ (inset in Fig. 5). However, a small change in the dc bias value E can tune the system into an instability window. As a result, the wave packet shows subdiffusive spreading on the accessible time scales, starting with $t \sim 10^5$ (inset in Fig. 5).




FIG. 5. (Color online) Amplitudes of the excited, $\nu=0$, and neighboring modes, $\nu=1$ and $\nu=2$, versus time for $\beta=0.4$. (a) $E=1.6$, $\nu=0$ (b, blue) and $E=2$, $\nu=0,1,2$ (o, orange; r, red; g, green). (b) $E=1.18$, $\nu=0$ (b, blue) and $E=1.26$, $\nu=0,1,2$ (o, orange; r, red; g, green). Inset: long time evolution. m_2 versus time in log-log plots: $E=1.18, 1.26$ (b, blue; o, orange).

These results can be observed experimentally for light propagating in waveguide networks [5]. Related instabilities of nonlinear Wannier-Stark states were briefly mentioned in [10].

In conclusion, we have investigated various regimes of wave-packet spreading for the nonlinear Stark ladder. For large nonlinearities there is a long-lived trapped regime with an explosive transition to Bloch oscillations, followed by a subdiffusive spreading at large time scales. For intermediate nonlinearities we find subdiffusion from scratch. For small nonlinearities the system is showing localization as a transient, with a subsequent subdiffusive spreading. For single-mode excitations additional stability and instability intervals with respect to the dc bias strength exist. For weak nonlinearity this leads to an additional sensitive dependence of the transient localization times on the value of the dc bias. In all cases, we observe that nonlinearity destroys integrability, introduces chaos, and ultimately leads to a destruction of localization, which was a hallmark of the linear system due to phase coherence and destructive and constructive wave interference.

The authors thank A. Ponomarev, Ch. Skokos, and U. Peschel for helpful discussions and comments. R.K. acknowledges financial support of the Georgian National Science Foundation (Grant No. GNSF/STO7/4-197).

- [1] R. Tsu, in *Semiconductor Interfaces, Microstructures and Devices*, edited by Z. C. Feng (IOP, Bristol, 1993).
 [2] G. H. Wannier, Phys. Rev. **117**, 432 (1960).
 [3] L. D. Landau, Phys. Z. Sowjetunion **2**, 46 (1932); G. Zener, Proc. R. Soc. London, Ser. A **137**, 696 (1932).
 [4] T. Pertsch, P. Dannberg, W. Elflein, A. Brauer, and F. Lederer, Phys. Rev. Lett. **83**, 4752 (1999); R. Sapienza, P. Costantino, D. Wiersma, M. Ghulinyan, C. J. Oton, and L. Pavesi, *ibid.* **91**, 263902 (2003); H. Trompeter, T. Pertsch, F. Lederer, D. Michaelis, U. Streppel, A. Brauer, and U. Peschel, *ibid.* **96**,

- 023901 (2006); F. Dreisow, A. Szameit, M. Heinrich, T. Pertsch, S. Nolte, A. Tunnermann, and S. Longhi, *ibid.* **102**, 076802 (2009).
 [5] R. Morandotti, U. Peschel, J. S. Aitchison, H. S. Eisenberg, and Y. Silberberg, Phys. Rev. Lett. **83**, 4756 (1999).
 [6] B. P. Anderson and M. A. Kasevich, Science **282**, 1686 (1998); M. Ben Dahan, E. Peik, J. Reichel, Y. Castin, and C. Salomon, Phys. Rev. Lett. **76**, 4508 (1996).
 [7] M. Gustavsson, E. Haller, M. J. Mark, J. G. Danzl, G. Rojas-Kopeinig, and H. C. Nagerl, Phys. Rev. Lett. **100**, 080404

- (2008).
- [8] O. Morsch, J. H. Muller, M. Cristiani, D. Ciampini, and E. Arimondo, *Phys. Rev. Lett.* **87**, 140402 (2001); G. Ferrari, N. Poli, F. Sorrentino, and G. M. Tino, *ibid.* **97**, 060402 (2006).
- [9] H. Fukuyama, R. A. Bari, and H. C. Fogedby, *Phys. Rev. B* **8**, 5579 (1973); P. Feuer, *Phys. Rev.* **88**, 92 (1952).
- [10] T. Pertsch, U. Peschel, and F. Lederer, *Phys. Rev. E* **66**, 066604 (2002).
- [11] P. K. Datta and A. M. Jayannavar, *Phys. Rev. B* **58**, 8170 (1998).
- [12] Y. Lahini, A. Avidan, F. Pozzi, M. Sorel, R. Morandotti, D. N. Christodoulides, and Y. Silberberg, *Phys. Rev. Lett.* **100**, 013906 (2008).
- [13] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (1972).
- [14] S. Flach, D. O. Krimer, and Ch. Skokos, *Phys. Rev. Lett.* **102**, 024101 (2009).
- [15] Ch. Skokos, D. O. Krimer, S. Komineas, and S. Flach, *Phys. Rev. E* **79**, 056211 (2009).
- [16] A. S. Pikovsky and D. L. Shepelyansky, *Phys. Rev. Lett.* **100**, 094101 (2008).
- [17] I. García-Mata and D. L. Shepelyansky, *Eur. Phys. J. B* (in print 2009).
- [18] A. Kolovsky, E. Gómez, and H. Korsch, e-print arXiv:0904.4549.
- [19] S. Raghavan, A. Smerzi, S. Fantoni, and S. R. Shenoy, *Phys. Rev. A* **59**, 620 (1999).
- [20] J. Laskar and P. Robutel, *Celest. Mech. Dyn. Astron.* **80**, 39 (2001).
- [21] Note, however, that T_E is sensitively depending on the integration time step in a fluctuating way.