

Formation of three-particle clusters in heterojunctions and MOSFET structures

E. P. Nakhmedov^{1,2} and K. Morawetz¹

¹Max-Planck-Institute for the Physics of Complex Systems, Noethnitzer Str. 38, 01187 Dresden, Germany

²Azerbaijan Academy of Sciences, Institute of Physics, H. Cavid 33, Baku, Azerbaijan

(Received 16 May 2002; revised manuscript received 13 September 2002; published 27 November 2002)

A novel interaction mechanism in metal-oxide-semiconduction field-effect transistor (MOSFET) structures and GaAs/AlGaAs heterojunctions between the zone electrons of a two-dimensional (2D) gas and charged traps on the insulator side is considered. By applying a canonical transformation, off-diagonal terms in the Hamiltonian due to the trapped level subsystem are excluded. This yields an effective three-particle attractive interaction as well as a pairing interaction inside the 2D electronic band. A type of Bethe-Goldstone equation for three particles is studied to clarify the character of the binding and the energy of the three-particle bound states. The results are used to offer a possible explanation of the metal-insulator transition recently observed in MOSFET and heterojunctions.

DOI: 10.1103/PhysRevB.66.195333

PACS number(s): 71.30.+h, 71.10.-w, 73.21.-b, 73.30.+y

Significant advances in low-temperature physics are particularly connected with the recent successes in fabricating two-dimensional (2D) electronic structures characterized by high mobility. The formation of the inversion layer of particles on interfaces allows the study of the unusual low-temperature behavior of a 2D electron gas, as well as the examination of theoretical predictions. Particularly, experimental measurements at very low temperatures $T \lesssim 2$ K show quite nonstandard results, like the fractional quantum Hall effect¹ and the metal-insulator transition (MIT),^{2,3} the origin of which is not yet properly understood. It is worthwhile to observe that the fractional quantum Hall effect and MIT occur only under some special regimes where the temperature and impurity concentrations in the samples are very small. On the other hand the observation of such unusual effects only in GaAs/AlGaAs heterojunctions and Si metal-oxide-semiconduction field-effect transistors (MOSFET) raises the question whether structural peculiarities of these devices are responsible or whether some fundamental law is observed.

The band structures of GaAs/AlGaAs heterojunctions and of Si-MOSFET's are well known, and their general features are identical despite their different structures. The inversion of current carriers occurs at the interface of two semiconductors (or insulator/semiconductor) with different band gaps. These semiconductors are exclusively doped by p - and n -type impurities in order to get high mobility in the samples. In extremely clean samples, the doping by one type of either acceptors or donors forms single levels in the gap of each semiconductor. In the process of, e.g., electron inversion, the donor centers in SiO₂ of the MOSFET structure (or in Al_xGa_{1-x}As heterojunctions) become positively charged by transferring electrons to the 2D electronic band. The charged donors are located within a region of at most ~ 200 Å (Ref. 4) from the oxide-silicon interface, and their energy level lies above the Fermi level. Such interfacial charged states act as trap centers for the band electrons. The density of trapped centers in, e.g., MOSFET structures is of the order of 10^9 cm⁻²,⁴ which is considerably smaller than the typical carrier concentration of $\sim 2 \times 10^{11}$ cm⁻². The Coulomb potential of charged traps seems to be screened

imperfectly due to their low densities, and the scattering of the band particles on these trap centers may be more essential than the intraband particle-particle scattering. In this paper we study the effects of the scattering of band electrons on the charged traps.

The Hamiltonian of the model can be written in the form $H = H_0 + H_{int}$, where

$$H_0 = \sum_{\mathbf{k}, \sigma} \varepsilon(\mathbf{k}) a_{\mathbf{k}, \sigma}^\dagger a_{\mathbf{k}, \sigma} + \sum_{\sigma} \omega_0 b_{0, \sigma}^\dagger b_{0, \sigma} \quad (1)$$

and

$$H_{int} = \sum_{\mathbf{k}, \mathbf{q}; \sigma, \sigma'} V(\mathbf{k}) [a_{\mathbf{k}, \sigma}^\dagger a_{-\mathbf{k}+\mathbf{q}, \sigma'}^\dagger b_{0, \sigma'} a_{\mathbf{q}, \sigma} + a_{\mathbf{q}, \sigma}^\dagger b_{0, \sigma'}^\dagger a_{-\mathbf{k}+\mathbf{q}, \sigma'} a_{\mathbf{k}, \sigma}]. \quad (2)$$

In Eqs. (1) and (2), $a_{\mathbf{k}, \sigma}^\dagger$ and $b_{0, \sigma}^\dagger$ ($a_{\mathbf{k}, \sigma}$ and $b_{0, \sigma}$) are the creation (annihilation) operators for the electrons band and in the trapped levels, respectively.

The trap centers are modeled for simplicity as a dispersionless single level. $\varepsilon(\mathbf{k})$ and ω_0 in Eq. (1) are the energies band electrons and trap level, respectively. The energy of the trap centers is considered to be larger than the chemical potential μ of the band electrons, $\omega_0 \geq \mu$. The first term in the Hamiltonian H_{int} represents the scattering of two band electrons via the interaction potential $V(\mathbf{k})$ followed by the trapping of one of them by the donor level. The second term represents the scattering of a band electron with a trapped electron (on a donor level) with finally turning both of them into the band.

For temperatures $kT < (\omega_0 - \mu)$, the trapping centers contain a definite number of electrons at thermodynamic equilibrium, and the considered trapping mechanism is assumed to be essential. However, the mechanism is destroyed by increasing the gate potential, for then the chemical potential μ reaches the trap level, causing the filling of all trapping centers. A temperature increase also leads to the destruction of the trapping mechanism.

We apply a unitary transformation⁵ with the intention to result in the cancellation of the off-diagonal term, given by Eq. (2). We will show here that this unitary transformation creates a “trap-mediated” effective attraction between three electrons in the band.⁶ Notice that our approach to the problem is similar to the cancellation of the phonon subsystem in superconductivity, where a canonical transformation yields an effective attraction between electrons; see, e.g., Ref. 7. Let us expand a new Hamiltonian $\tilde{H} = e^{-iS} H e^{iS}$ in a power series of the operator $S = S_1 + S_2 + S_3 + \dots$. Regrouping the terms of the same order in H_{int} , the conditions which define $S_i, i = 1, 2, \dots$ are obtained recursively. S_1 is determined by $H_{int} - i[S_1, H_0] = 0$ and leads to

$$S_1 = i \sum_{\mathbf{k}, \mathbf{q}; \sigma, \sigma'} \frac{V_{\mathbf{k}}}{\epsilon_{\mathbf{k}} + \epsilon_{-\mathbf{k}+\mathbf{q}} - \epsilon_{\mathbf{q}} - \omega_0} [a_{\mathbf{k}, \sigma}^\dagger a_{-\mathbf{k}+\mathbf{q}, \sigma'}^\dagger b_{0, \sigma'} a_{\mathbf{q}, \sigma} - a_{\mathbf{q}, \sigma}^\dagger b_{0, \sigma'}^\dagger a_{-\mathbf{k}+\mathbf{q}, \sigma'} a_{\mathbf{k}, \sigma}]. \quad (3)$$

The new Hamiltonian \tilde{H} can now be written in the form $\tilde{H} = H_0 - (i/2)[S_1, H_{int}] - i[S_2, H_0]$. Here S_2 is obtained from the condition that the off-diagonal terms in the Hamiltonian be cancelled. The equation for \tilde{H} then becomes $\tilde{H} = H_0 - (i/2)[S_1, H_{int}]_{diag}$, where the last term contains only diagonal elements. Introducing Eq. (3) into the expression for \tilde{H} finally yields the effective Hamiltonian

$$\tilde{H} = \tilde{H}_0 + H_{e-e} + H_{\Delta}, \quad (4)$$

where \tilde{H}_0 , H_{e-e} , and H_{Δ} describe the effective one-particle Hamiltonian, the electron-electron interaction, and the three-particle clustering, respectively. The one-particle effective Hamiltonian \tilde{H}_0 is given by

$$\begin{aligned} \tilde{H}_0 = & H_0 - \sum_{\sigma'} b_{0, \sigma'}^\dagger b_{0, \sigma'} \sum_{\mathbf{q}, \sigma} \varepsilon_1(\mathbf{q}) a_{\mathbf{q}, \sigma}^\dagger a_{\mathbf{q}, \sigma} \\ & + \sum_{\mathbf{q}, \sigma, \sigma'} J(\mathbf{q}) b_{0, \sigma}^\dagger b_{0, \sigma'}^\dagger a_{\mathbf{q}, \sigma'}^\dagger a_{\mathbf{q}, \sigma}, \\ \left(\begin{array}{c} \varepsilon_1(\mathbf{q}) \\ J(\mathbf{q}) \end{array} \right) = & \sum_{\mathbf{k}} \frac{1}{\epsilon_{\mathbf{k}} - \epsilon_{-\mathbf{k}+\mathbf{q}} - \epsilon_{\mathbf{q}} - \omega_0} \left(\begin{array}{c} V_{\mathbf{k}}^2 \\ V_{\mathbf{k}} V_{-\mathbf{k}+\mathbf{q}} \end{array} \right). \end{aligned} \quad (5)$$

In thermodynamical equilibrium $\langle b_{0, \sigma}^\dagger, b_{0, \sigma} \rangle = \delta_{\sigma, \sigma'}$, and the one-particle energy is renormalized, $\tilde{H}_0 = \sum_{\mathbf{q}, \sigma} \tilde{\varepsilon}(\mathbf{q}) a_{\mathbf{q}, \sigma}^\dagger a_{\mathbf{q}, \sigma}$, with $\tilde{\varepsilon}(\mathbf{q}) = \varepsilon(\mathbf{q}) + \varepsilon_1(\mathbf{q}) + J(\mathbf{q})$.

The electron-electron interaction Hamiltonian H_{e-e} also contains terms with spin flipping due to the exchange scattering of 2D electrons on trapped ones. In thermodynamic equilibrium, H_{e-e} has the usual form

$$H_{e-e} = \frac{1}{2} \sum_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}; \sigma, \sigma'} V_{e-e}^{(eff)}(\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}) a_{\mathbf{k}_1, \sigma}^\dagger a_{\mathbf{k}_2, \sigma'}^\dagger a_{\mathbf{k}_2 - \mathbf{q}, \sigma'} a_{\mathbf{k}_1 + \mathbf{q}, \sigma}, \quad (6)$$

where the effective two-particle interaction potential $V_{e-e}^{(eff)}(\mathbf{k}_1, \mathbf{k}_2, \mathbf{q})$ appears to be attractive,

$$\begin{aligned} V_{e-e}^{(eff)}(\mathbf{k}_1, \mathbf{k}_2, \mathbf{q}) = & \frac{V_{\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{q}}^2}{\epsilon_{\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{q}} + \epsilon_{\mathbf{k}_2} - \epsilon_{\mathbf{k}_1 + \mathbf{q}} - \omega_0} \\ & - \frac{V_{\mathbf{k}_1} V_{\mathbf{k}_2 - \mathbf{q}}}{\epsilon_{\mathbf{k}_1} + \epsilon_{\mathbf{q}} - \epsilon_{\mathbf{k}_1 + \mathbf{q}} - \omega_0} \\ & - \frac{V_{\mathbf{k}_1} V_{\mathbf{k}_2 - \mathbf{q}}}{\epsilon_{\mathbf{k}_2 - \mathbf{q}} + \epsilon_{\mathbf{q}} - \epsilon_{\mathbf{k}_2} - \omega_0} \\ & + \frac{V_{\mathbf{k}_1} V_{\mathbf{q}} + V_{\mathbf{k}_2 - \mathbf{q}} V_{\mathbf{q}}}{\epsilon_{\mathbf{q}} + \epsilon_{\mathbf{k}_2 - \mathbf{q}} - \epsilon_{\mathbf{k}_2} - \omega_0} \\ & + \frac{V_{\mathbf{k}_1} V_{\mathbf{q}} + V_{\mathbf{k}_2 - \mathbf{q}} V_{\mathbf{q}}}{\epsilon_{\mathbf{q}} + \epsilon_{\mathbf{k}_1} - \epsilon_{\mathbf{k}_1 + \mathbf{q}} - \omega_0} \\ & + \frac{V_{\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{q}}^2}{\epsilon_{\mathbf{k}_1 - \mathbf{k}_2 + \mathbf{q}} + \epsilon_{\mathbf{k}_2 - \mathbf{q}} - \epsilon_{\mathbf{k}_1} - \omega_0}. \end{aligned} \quad (7)$$

Indeed, the denominator of each term in $V_{e-e}^{(eff)}$ is negative, owing to the fact that a donor level lies higher than the chemical potential of band electrons.

The third term in the Hamiltonian (4) describes an effective three-particle scattering,

$$\begin{aligned} H_{\Delta} = & \frac{1}{2} \sum_{\substack{\sigma_1, \sigma_2, \sigma_3 \\ \mathbf{k}_1, \mathbf{k}_2, \mathbf{q}_1, \mathbf{q}_2}} \left(\frac{V_{\mathbf{k}_1} V_{\mathbf{k}_2}}{\epsilon_{\mathbf{k}_1} + \epsilon_{-\mathbf{k}_1 + \mathbf{q}_1} - \epsilon_{\mathbf{q}_1} - \omega_0} \right. \\ & \left. + \frac{V_{\mathbf{k}_1} V_{\mathbf{k}_2}}{\epsilon_{\mathbf{k}_2} + \epsilon_{-\mathbf{k}_2 + \mathbf{q}_2} - \epsilon_{\mathbf{q}_2} - \omega_0} \right) a_{\mathbf{q}_1, \sigma_1}^\dagger a_{\mathbf{k}_2, \sigma_2}^\dagger a_{-\mathbf{k}_2 + \mathbf{q}_2, \sigma_3} \\ & a_{-\mathbf{k}_1 + \mathbf{q}_1, \sigma_3} a_{\mathbf{q}_2, \sigma_2} a_{\mathbf{k}_1, \sigma_1}. \end{aligned} \quad (8)$$

Using again the condition $\mu < \omega_0$, it is possible to see that the strength of the three-particle interaction is negative, which results in the formation of clusters of three electrons.

This effective attraction among three electrons can be understood according to the following physical argument. The proposed mechanism of two-particle interactions with trapping of one of the particles, in contrast to an intraband electron-electron scattering, destroys locally the electroneutrality of the 2D electron gas. The necessary electroneutrality in heterojunctions or in MOSFET's is restored by the ensuing adaption of the height of the Schottky barrier, i.e., by a change in the value of the band bending energy. However, the trapping and releasing processes are so fast that the barrier's height cannot follow. As a result of the trapping of band electrons, a hole appears which acts as an attractive center for other electrons.

The energy level of the trap centers in the above calculation is chosen to be dispersionless for simplicity. However, even in the single-level case, the donor center energies depend on the spatial coordinates of the impurities due to the band bending and, therefore, become dispersive. Including the dispersion of the trap level does not change qualitatively our results.

We now proceed to show that a three-particle attractive interaction can lead to the formation of a bound state. To this end, we consider for simplicity only the three-particle interaction and neglect the pairing interaction. The Schrödinger equation for three identical particles in 2D with a generic interaction potential of the form $V(\mathbf{r}_1 - \mathbf{r}_2; \mathbf{r}_3 - \mathbf{r}_1; \mathbf{r}_2 - \mathbf{r}_3)$ is written in the following form after introducing the Jacobi coordinates $\mathbf{R} = \frac{1}{3}(\mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3)$, $\mathbf{r} = \mathbf{r}_3 - \mathbf{r}_1$, and $\mathbf{z} = (\mathbf{r}_1 + \mathbf{r}_3)/2 - \mathbf{r}_2$:

$$\left\{ -\frac{\hbar^2}{2m} \left(\frac{1}{3} \frac{\partial^2}{\partial \mathbf{R}^2} + 2 \frac{\partial^2}{\partial \mathbf{r}^2} + \frac{2}{3} \frac{\partial^2}{\partial \mathbf{z}^2} \right) + V \left(\mathbf{z} - \frac{\mathbf{r}}{2}; \mathbf{r}; \mathbf{z} + \frac{\mathbf{r}}{2} \right) \right\} \psi(\mathbf{R}, \mathbf{r}, \mathbf{z}) = (\epsilon + 3\epsilon_F) \psi(\mathbf{R}, \mathbf{r}, \mathbf{z}), \quad (9)$$

where ϵ is the three-particle excitation energy measured from the three-particle Fermi level. After excluding the center-of-mass coordinate \mathbf{R} by expanding $\psi(\mathbf{R}, \mathbf{r}, \mathbf{z})$ in plane waves, $\psi(\mathbf{R}, \mathbf{r}, \mathbf{z}) = \sum_{\mathbf{Q}, \mathbf{p}, \mathbf{q}} e^{(i/\hbar)(\mathbf{Q}\mathbf{R} + \mathbf{p}\mathbf{r} + \mathbf{q}\mathbf{z})} \phi(\mathbf{p}, \mathbf{q})$, a Bethe-Goldstone-type equation, similar to the equation for Cooper pairs,⁸ is obtained,

$$\left(\frac{\mathbf{p}^2}{m} + \frac{3\mathbf{q}^2}{4m} - \epsilon - 3\epsilon_F \right) \phi(\mathbf{p}, \mathbf{q}) + \sum_{\mathbf{p}', \mathbf{q}'} \tilde{V}(\mathbf{p}', \mathbf{q}'; \mathbf{p}, \mathbf{q}) \phi(\mathbf{p}', \mathbf{q}') = 0. \quad (10)$$

The interaction potential $\tilde{V}(\mathbf{p}', \mathbf{q}'; \mathbf{p}, \mathbf{q})$ is assumed to be attractive when the energies of the three particles (before the coordinate transformation) lie in a narrow vicinity $\hbar\omega_0$ of the Fermi surface ($\epsilon_F, \epsilon_F + \hbar\omega_0$). Here $\hbar\omega_0$ is a cutoff energy which is comparable to the order of trap level energy measured from the Fermi level. This condition restricts the energies $\frac{3}{4}\epsilon_F < p^2/2m < \frac{3}{4}(\epsilon_F + \hbar\omega_0)$ and $\epsilon_F < q^2/2m < \epsilon_F + \hbar\omega_0$ of the quasiparticles obtained after coordinate transformation. Therefore, the simplified attractive interaction for a system of linear size L is $\tilde{V}(\mathbf{p}', \mathbf{q}'; \mathbf{p}, \mathbf{q}) = -V_0/L^4$ for $\frac{3}{4}\epsilon_F < p^2/2m$, $\frac{p'^2}{2m} < \frac{3}{4}(\epsilon_F + \hbar\omega_0)$, $\epsilon_F < \frac{q^2}{2m}$, $\frac{q'^2}{2m} < \epsilon_F + \hbar\omega_0$, and 0 otherwise.

The following equation for the bound-state energy of a three-particle cluster results,

$$1 = \frac{1}{3} V_0 N^2 \int_0^{3\hbar\omega_0/2} d\xi_1 \int_0^{3\hbar\omega_0/2} d\xi_2 \frac{1}{\xi_1 + \xi_2 - \epsilon}, \quad (11)$$

where N is the value of the density of electronic states on the Fermi surface. The integration of Eq. (11) gives an equation for ϵ ,

$$\frac{3}{V_0 N^2 \hbar\omega_0} = 3 \ln \frac{6 - 2\tilde{\epsilon}}{3 - 2\tilde{\epsilon}} - \tilde{\epsilon} \ln \frac{-4\tilde{\epsilon}(3 - \tilde{\epsilon})}{(3 - 2\tilde{\epsilon})^2}, \quad (12)$$

where $\tilde{\epsilon} = \epsilon/\hbar\omega_0$ is the dimensionless excitation energy. The study of Eq. (12) for arbitrary negative values of $\tilde{\epsilon}$ shows that the right-hand side of this equation is a monotonic and positive function with a maximum value equal to $\ln 8$ at $\tilde{\epsilon} = 0$. This imposes a lower restriction on the attractive potential, $V_0 \geq 1/\ln 2 N^2 \hbar\omega_0$. Consequently, for attractive potentials strong enough, Eq. (12) possesses a unique negative solution for $-\tilde{\epsilon} \in (-\infty, 0)$. This implies the existence of three-particle bound states.

In a weak-coupling regime, when $-\tilde{\epsilon} \ll 1$, Eq. (12) is simplified to

$$\frac{-4\tilde{\epsilon}}{3e} \ln \frac{-4\tilde{\epsilon}}{3e} = \frac{4}{e} \left(\frac{1}{V_0 N^2 \hbar\omega_0} - \ln 2 \right), \quad (13)$$

the solution of which does not possess a gaplike form for the excitation energy.

In the opposite case of large negative solutions, $-\tilde{\epsilon} \geq 3$, Eq. (12) leads immediately to the following result for the bound energy in the strong-coupling regime, $\epsilon = -\frac{3}{4} V_0 (\hbar\omega_0 N)^2$, which shows a clear perturbative noncollective behavior.

This discussion shows that molecular clustering rather than a coherent state is realized in the system. The ground state of the system becomes unstable with respect to the three-particle attraction. This seems to lead to molecular-type formation with negative energy.

In conclusion, we want to emphasize that it is possible to understand qualitatively the reason of MIT's occurring at very low temperatures in GaAs/AlGaAs heterojunctions and MOSFET (Ref. 2 and 9) in the framework of the formation of three-particle bound states we describe above. The elastic scattering of electrons on impurities at low temperatures, which is characterized by a relaxation time τ_0 , results in the localization of all electronic states,¹⁰ under the condition that $\hbar\omega_0(n) < kT < \hbar/\tau_0$, producing an insulating behavior for conductivity. Here $\omega_0(n)$ is the trap level energy measured from the Fermi energy, which is a function of the 2D electron concentration n , or the gate potential. On the other hand, in a regime corresponding to $kT < \hbar\omega_0(n) < \hbar/\tau_0$, which can be reached by varying the electron concentration or the temperature, the formation of three-particle bound states results in the vanishing of the weak-localization corrections to conductivity. This is due to the fact that the scattering of the three-particle clusters on the impurities does not lead to quantum interference. Instead, the cluster's wave function accumulates an additional phase by rotation of the cluster in the process of scattering, while the center-of-mass motion of the cluster is still extended. The expression for the conductivity can be written as

$$\sigma(T) = \frac{3e^2 n \Delta \tau_0}{m} + \frac{e^2 n_f \tau_0}{m} \left(1 - \frac{\hbar}{2\pi \epsilon_F \tau_0} \ln \frac{\tau_{in}(T)}{\tau_0} \right), \quad (14)$$

where the first and second terms in Eq. (14) correspond to the Drude and weak-localization contributions,¹¹ which correspond to three-particles clusters and free band electrons




FIG. 1. The temperature dependence of the experimental (dots) conductivity scaled to the critical one (Ref. 12). The densities increase from below to above as $n = 6.85, 7.17, 7.25, 7.57, 7.85 \times 10^{10} \text{ cm}^{-2}$. The lines are theoretical fits according to Eq. (15) where the lower three curves are assumed to be undercritical (without n_Δ) with the fit $T_c = 3.9, 3.6, 1.7 \text{ K}$ and $\lambda = 0.330, 0.078, 0.004$ correspondingly and the above two metallic curves are critical (with n_Δ) with the fit $T_c = 1.4, 0.8 \text{ K}$ and $\lambda = 0.48, 0.86$.

with concentrations of n_Δ and n_f , respectively. τ_{in} is the inelastic scattering time $\tau_{in} = aT^{-p}$ where a is some constant, $p \geq 2$ and $p = 2$ for a probable electron-electron scattering mechanism. Notice that a logarithmic temperature dependence of σ in the metallic phase has been observed³ in high-mobility n -Si-MOSFET's which is in good agreement with our assumption. Observation of the negative low-field magnetoresistance in the metallic phase also supports the important role of quantum interference effects in MIT's. We neglect in Eq. (14) an additional logarithmic quantum correction due to the electron-electron interactions,¹¹ which is responsible for the positive magnetoresistance also observed in experiments.²

Charge conservation allows us to write the total concentration of particles as $n = n_\Delta + n_f + n_t$, where n_t is the concentration of trapped electrons, which exponentially decreases with temperature as $n_t(T) = n_t^0 \exp\{-\omega(n)/kT\}$ with n_t^0 being the concentration of trapped impurities. Assuming that clustering occurs at $T = T_c(n)$, n_Δ can be expressed near $T_c(n)$ as $n_\Delta/n = (T_c - T)/T_c$. We rewrite Eq. (14) in the form

$$\frac{\sigma(T)}{\sigma_0} = 1 + 2 \frac{T_c - T}{T_c} - \frac{\hbar T}{\pi \epsilon_F \tau_0 T_c} \ln \frac{T^*}{T}, \quad (15)$$

where $\sigma_0 = e^2 \tau_0 n / m$ is the Drude conductivity, and $T^* = \sqrt{a} / \tau_0$. In Eq. (15) we neglected the trap level contribution due to $n_t \ll n$. Rescaling T by T^* as $T/T^* \equiv \tau$ for $\tau < 1$ and choosing the parameter of randomness $\lambda = \hbar / 2\pi \epsilon_F \tau_0 \leq 1$, the unknown parameter $T_c/T^* \equiv \tilde{T}_c$ in Eq. (15) can be extracted by fitting $\sigma(T)/\sigma_0$ to the experimental data. The temperature dependence of σ is drawn in Fig. 1 for the best fit parameters $\tilde{T}_c(n)$.

While the overall fit to MIT's is satisfactorily we see deviations on the insulating side. This seems to be due to

strong-localization effects in these experiments. Effects observed in the weak-localization regime³ show a more pronounced logarithmic behavior. The values of the conductivity at the critical density G_c observed in the two experiments strongly differ from each other: in the strongly disordered case,² $G_c \sim e^2/h$, whereas in the weakly disordered case,³ $G_c \sim 120e^2/h$. Therefore, we think that the conductivity at the critical density G_c does not show a universal behavior. It depends on two factors: on the impurity concentration in the Si substrate or in the 2D electron gas and on the electron concentration (or on the band filling) in the inversion layer. By increasing the band filling in the insulator side of the MIT's three-particle clusters appear which weaken the localization tendency in the 2D electronic system since the scattering off the cluster on the impurities does not lead to quantum interference effects. At the critical density, the contributions coming from the clustering completely compensate the localization corrections and the conductivity is defined by the value of the residual Drude conductivity, which is temperature independent at low temperature.

Notice that another mechanism for MIT's, which is controlled by a temperature-dependent trapped-electron concentration $n_t(T)$, has been recently proposed by Altshuler and Maslov.¹³ As the Comment and Reply show,¹⁴ this qualitatively correct explanation cannot reproduce quantitative features of the experiment. A critical discussion of different approaches can be found in Ref. 2.

Although we have not discussed a role of effective pairing governed by Eq. (6) in the Hamiltonian, there was an attempt to interpret the experimental data on MIT's as a result of possible superconducting ground state.¹⁵ It is well known that the effective pairing is suppressed by the order parameter phase fluctuations in 2D systems reducing T_{SC} of the superconducting transition to zero. However, fluctuations of the order parameter modulus above T_{SC} may lead to the metallic phase.

The geometry of the cluster may be either in the form of a triangle with 3/2 and 1/2 total spin or of a stringlike configuration with 1/2 total spin, when two electrons with antiparallel spins are placed at the same point and the third electron with arbitrary spin is far from them. In the case of 3/2 total spin, a magnetic field parallel to the triangle area does not destroy the cluster, whereas the configurations with 1/2 total spin are destroyed due to the Zeeman effect. In both cases the magnetic field effects are defined by the contributions coming from the quantum localization corrections.

The model of three-particle clustering due to the discussed exchange type of interaction with donor levels seems to be also a favorable candidate for the understanding of the fractional quantum Hall effect. The interaction of the band electrons with trap centers effectively leads to the formation of three-particle clusters [see Eq. (8)], as well as to the superconducting fluctuations due to effective pairing interactions [Eq. (6)]. Both mechanisms decrease the ground-state energy of the system. Strong magnetic fields in the quantum

Hall regime polarize the spins of molecular clusters and a triangular geometry for the cluster is realized due to the Pauli principle. An antisymmetric orbital wave function of the triangular cluster will contain *ab initio* the Jastrow prefactor. The angular momentum $M = 3$ of the cluster provides a natu-

ral argument in Laughlin's theory to connect the filling factor $\nu = 1/3$ of the parent states with the angular momentum $M = 3$.

The authors gratefully acknowledge discussions with M. Ameduri, D. Efremov, P. Fulde, and K. Maki.

-
- ¹R.E. Prange and S.M. Girvin, *The Quantum Hall Effect* (Springer-Verlag, Berlin, 1987).
- ²E. Abrahams, S.V. Kravchenko, and M.P. Sarachik, *Rev. Mod. Phys.* **73**, 251 (2001).
- ³V. Pudalov *et al.*, *JETP Lett.* **68**, 534 (1998).
- ⁴R.M. Warner and J.B.L. Grung, *MOSFET: Theory and Design* (Oxford University Press, Oxford, 1999).
- ⁵J.R. Schrieffer and P.A. Wolff, *Phys. Rev.* **149**, 491 (1966).
- ⁶E.P. Nakhmedov, *Turk. J. Phys.* **21**, 155 (1997).
- ⁷C. Kittel, *Quantum Theory of Solids* (Wiley, New York, 1987).
- ⁸P. de Gennes, *Superconductivity of Metals and Alloys* (Addison-Wesley, New York, 1989).
- ⁹S.V. Kravchenko *et al.*, *Phys. Rev. B* **50**, 8039 (1994).
- ¹⁰E. Abrahams, P.W. Anderson, D.C. Licciardello, and T.V. Ramakrishnan, *Phys. Rev. Lett.* **42**, 673 (1979).
- ¹¹P.A. Lee and T.V. Ramakrishnan, *Rev. Mod. Phys.* **57**, 287 (1985).
- ¹²S.V. Kravchenko and T.M. Klapwijk, *Phys. Rev. Lett.* **84**, 2909 (2000).
- ¹³B.L. Altshuler and D.L. Maslov, *Phys. Rev. Lett.* **82**, 145 (1999).
- ¹⁴S.V. Kravchenko, M.P. Sarachik, and D. Simonian, *Phys. Rev. Lett.* **83**, 2091 (1999).
- ¹⁵P. Phillips *et al.*, *Nature (London)* **395**, 253 (1998).